

# **Supplemental Material**

## Appendix

### The International Collaboration of Comprehensive Physiologic Assessment Investigators

Name	Affiliations
Seung Hun Lee, MD, PhD	Division of Cardiology, Department of Internal Medicine, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea
Joo Myung Lee, MD, MPH, PhD	Division of Cardiology, Department of Internal Medicine, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea
Jonghanne Park, MD, PhD	Department of Internal Medicine and Cardiovascular Center, Seoul National University Hospital, Seoul, Korea Developmental Therapeutics Program of Division of Hematology Oncology, Northwestern University, Chicago, Illinois, USA
Ki Hong Choi, MD	Division of Cardiology, Department of Internal Medicine, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea
Doyeon Hwang, MD	Department of Internal Medicine and Cardiovascular Center, Seoul National University Hospital, Seoul, Korea
Joon-Hyung Doh, MD, PhD	Department of Medicine, Inje University Ilsan Paik Hospital, Goyang, Korea
Chang-Wook Nam, MD, PhD	Department of Medicine, Keimyung University Dongsan Medical Center, Daegu, Korea
Eun-Seok Shin, MD, PhD	Division of Cardiology, Ulsan Hospital, Ulsan, Korea Department of Cardiology, Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, Korea
Ji-Hyun Jung, MD	Sejong General Hospital, Sejong Heart Institute, Bucheon, Korea
Hyung Yoon Kim, MD	Department of Cardiovascular Medicine, Chonnam National University Hospital, Gwangju, Korea
Hae Won Jung, MD	Department of Cardiology, Daegu Catholic University Medical Center, Daegu, Korea
Yun-Kyeong Cho, MD, PhD	Department of Medicine, Keimyung University Dongsan Medical Center, Daegu, South Korea
Hyuck-Jun Yoon, MD, PhD	Department of Medicine, Keimyung University Dongsan Medical Center, Daegu, South Korea
Young Bin Song, MD, PhD	Division of Cardiology, Department of Internal Medicine, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea
Joo-Yong Hahn, MD, PhD	Division of Cardiology, Department of Internal Medicine, Heart Vascular Stroke Institute, Samsung Medical Center,

Seung-Ho Hur, MD, PhD  
Masahiro Hoshino, MD  
Yoshihisa Kanaji, MD  
Rikuta Hamaya, MD  
Yoshinori Kanno, MD  
Masahiro Hada, MD  
Masao Yamaguchi, MD  
Hiroaki Ohya, MD  
Yohei Sumino, MD  
Eisuke Usui, MD  
Akinori Sugano, MD  
Tomoyo Sugiyama, MD, PhD  
Tadashi Murai, MD, PhD  
Tetsumin Lee, MD  
Taishi Yonetsu, MD  
Hernán Mejía-Rentería, MD  
Francesco Lauri, MD  
Sonoka Goto, MD  
Fernando Macaya, MD  
Angela McInerney, MD  
Giacomo Gravina, MD  
Rafael Vera, MD  
Nieves Gonzalo, MD  
Pilar Jimenez-Quevedo, MD  
Ivan Nuñez-Gil, MD  
Pablo Salinas, MD  
Luis Nombela-Franco, MD  
Maria del Trigo, MD  
Antonio Fernández-Ortiz, MD  
Carlos Macaya, MD  
Tsunekazu Kakuta, MD, PHD

Sungkyunkwan University School of Medicine, Seoul, Korea  
Department of Medicine, Keimyung University Dongsan Medical Center, Daegu, South Korea  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan  
Department of Cardiovascular Medicine, Tokyo Medical and Dental University, Tokyo, Japan  
Department of Cardiovascular Medicine, Tokyo Medical and Dental University, Tokyo, Japan  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Division of Cardiovascular Medicine, Tsuchiura Kyodo General Hospital, Ibaraki, Japan

Javier Escaned, MD, PhD

Bon-Kwon Koo, MD, PhD

Cardiovascular Institute, Hospital Clinico San Carlos, Madrid, Spain  
Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), Madrid, Spain  
Department of Internal Medicine and Cardiovascular Center, Seoul National University Hospital, Seoul, Korea  
Institute on Aging, Seoul National University, Seoul, Korea

---

**Table S1. Characteristics of Patients According to the Registries.**

	Total population	Korea registry	Japan registry	Spain registry
<b>Per-patient analysis</b>	<b>n=1,245</b>	<b>n=552</b>	<b>n=631</b>	<b>n=62</b>
<b>General characteristics</b>				
Age, years	64.7 ± 10.3	61.6 ± 10.3	67.4 ± 9.4	65.6 ± 10.8
Male	958 (76.9%)	389 (70.5%)	520 (82.4%)	49 (79.0%)
BMI, kg/m <sup>2</sup>	24.8 ± 3.5	24.6 ± 3.0	24.8 ± 3.8	28.5 ± 4.4
<b>Clinical presentation</b>				
Stable ischemic heart disease	1115 (89.6%)	454 (88.2%)	631 (100%)	30 (48.4%)
Acute coronary syndrome	130 (10.4%)	98 (17.8%)	0 (0%)	32 (51.6%)
<b>Cardiovascular risk factors</b>				
Hypertension	817 (65.6%)	328 (59.4%)	444 (70.4%)	45 (72.6%)
Diabetes mellitus	445 (35.7%)	160 (29.0%)	260 (41.2%)	25 (40.3%)
Hypercholesterolemia	784 (63.0%)	350 (63.4%)	400 (63.4%)	34 (54.8%)
Current smoker	266 (21.4%)	105 (19.0%)	142 (22.5%)	19 (30.6%)
Obesity (BMI>25 kg/m <sup>2</sup> )	548 (44.8%)	234 (42.7%)	280 (44.4%)	34 (79.1%)
Multivessel disease	435 (34.9%)	273 (49.5%)	113 (17.9%)	49 (79.0%)
<b>Per-vessel analysis</b>	<b>n=1,484</b>	<b>n=772</b>	<b>n=631</b>	<b>n=81</b>
<b>Vessel location</b>				
LAD	950 (64.0%)	463 (60.0%)	442 (70.1%)	45 (55.6%)
LCX	219 (14.8%)	136 (17.6%)	67 (10.6%)	16 (19.7%)

RCA	315 (21.2%)	173 (22.4%)	122 (19.3%)	20 (24.7%)
Target vessel PCI performed	464 (31.3%)	96 (12.4%)	338 (53.6%)	30 (37.0%)
<b>Angiographic characteristics</b>				
Reference diameter	2.9 ± 0.6	3.0 ± 0.6	2.8 ± 0.7	2.8 ± 0.7
Diameter stenosis, %	46.0 ± 16.8	42.5 ± 17.5	49.7 ± 15.2	53.2 ± 12.5
Lesion length, mm	12.6 ± 8.6	11.6 ± 8.3	12.6 ± 7.5	23.0 ± 13.0
<b>Physiologic parameters</b>				
Resting Pd/Pa	0.93 ± 0.08	0.95 ± 0.06	0.90 ± 0.09	0.89 ± 0.09
Fractional flow reserve	0.83 ± 0.11	0.86 ± 0.10	0.79 ± 0.11	0.80 ± 0.10
Coronary flow reserve	2.9 ± 1.3	3.1 ± 1.3	2.8 ± 1.3	2.1 ± 1.4
IMR, U	21.0 ± 13.9	19.4 ± 9.7	22.9 ± 17.1	20.9 ± 18.5
Resistive reserve ratio	3.7 ± 1.7	3.8 ± 1.6	3.6 ± 1.7	2.7 ± 1.9

---

Values expressed as mean ± SD or number (%).

BMI, body mass index; IMR, index of microcirculatory resistance; LAD, left anterior descending artery; LCX, left circumflex artery; Pa, aortic pressure; Pd, distal coronary pressure; RCA, right coronary artery; Tmn, mean transit time.

**Table S2. Characteristics of Patients with Deferred Revascularization According to Resistive Reserve Ratio in High CFR or High FFR Subgroups.**

	High FFR Population			High CFR Population		
	RRR $\geq$ 3.5	RRR<3.5	P value	RRR $\geq$ 3.5	RRR<3.5	P value
<b>Per-patient analysis (n=787)</b>	<b>349/643 (54.3%)</b>	<b>294/643 (45.7%)</b>		<b>416/597 (69.7%)</b>	<b>181/597 (30.3%)</b>	
<b>General characteristics</b>						
Age, years	62.4 $\pm$ 10.5	65.8 $\pm$ 10.0	<0.001	62.3 $\pm$ 10.3	65.0 $\pm$ 10.2	0.003
Male	268 (76.8%)	193 (65.7%)	0.002	331 (79.6%)	118 (65.2%)	<0.001
BMI, kg/m <sup>2</sup>	24.8 $\pm$ 3.9	24.7 $\pm$ 3.5	0.911	24.9 $\pm$ 3.8	24.6 $\pm$ 3.5	0.476
<b>Clinical presentation</b>			<b>0.827</b>			<b>1.000</b>
Stable ischemic heart disease	308 (88.2%)	262 (89.1%)		366 (88.0%)	159 (87.9%)	
Acute coronary syndrome	41 (11.8%)	32 (10.9%)		50 (12.0%)	22 (12.1%)	
<b>Cardiovascular risk factors</b>						
Hypertension	213 (61.0%)	187 (63.6%)	0.556	262 (63.0%)	109 (60.2%)	0.584
Diabetes mellitus	99 (28.4%)	115 (39.1%)	0.005	114 (27.4%)	74 (40.9%)	0.002
Hypercholesterolemia	219 (62.8%)	170 (57.8%)	0.233	263 (63.2%)	114 (63.0%)	1.000
Current smoker	74 (21.2%)	64 (21.8%)	0.938	87 (20.9%)	41 (22.7%)	0.713
Obesity (BMI>25 kg/m <sup>2</sup> )	154 (44.4%)	126 (43.9%)	0.968	189 (45.8%)	79 (44.9%)	0.916
Multivessel disease	120 (34.4%)	129 (43.9%)	0.017	155 (37.3%)	73 (40.3%)	0.536
<b>Per-vessel analysis (n=1,020)</b>	<b>470/866 (54.3%)</b>	<b>396/866 (45.7%)</b>		<b>540/767 (70.4%)</b>	<b>227/767 (29.6%)</b>	
<b>Vessel location</b>			<b>0.071</b>			<b>0.132</b>


LAD	261 (55.5%)	231 (58.3%)		324 (60.0%)	144 (63.4%)	
LCX	77 (16.4%)	79 (20.0%)		79 (14.6%)	40 (17.6%)	
RCA	132 (28.1%)	86 (21.7%)		137 (25.4%)	43 (18.9%)	
<b>Angiographic characteristics</b>						
Reference diameter	3.1 ± 0.6	2.9 ± 0.6	0.001	3.0 ± 0.6	2.9 ± 0.7	0.007
Diameter stenosis, %	39.0 ± 14.9	39.6 ± 14.8	0.501	40.2 ± 15.0	39.9 ± 14.6	0.789
Lesion length, mm	10.9 ± 7.0	10.9 ± 6.9	0.988	11.3 ± 7.6	10.5 ± 6.3	0.126
<b>Physiologic parameters</b>						
Resting Pd/Pa	0.96 ± 0.04	0.96 ± 0.04	<0.001	0.96 ± 0.04	0.95 ± 0.04	0.141
Fractional flow reserve	0.90 ± 0.05	0.90 ± 0.05	0.757	0.88 ± 0.07	0.88 ± 0.07	0.383
Coronary flow reserve	4.1 ± 0.9	2.0 ± 0.6	<0.001	4.0 ± 0.9	2.5 ± 0.3	<0.001
Resting Tmn, sec	0.99 ± 0.44	0.62 ± 0.40	<0.001	0.98 ± 0.43	0.72 ± 0.38	<0.001
Hyperemic Tmn, sec	0.25 ± 0.11	0.31 ± 0.21	<0.001	0.25 ± 0.11	0.29 ± 0.16	<0.001
IMR, U	18.9 ± 8.7	24.0 ± 18.3	<0.001	18.3 ± 8.5	22.3 ± 12.8	<0.001
Resistive reserve ratio	5.1 ± 1.2	2.4 ± 0.7	<0.001	5.1 ± 1.2	3.0 ± 0.3	<0.001

Values expressed as mean ± SD or number (%).

BMI, body mass index; IMR, index of microcirculatory resistance; LAD, left anterior descending artery; LCX, left circumflex artery; Pa, aortic pressure; Pd, distal coronary pressure; RCA, right coronary artery; RRR, resistive reserve ratio; Tmn, mean transit time.




**Figure S1. Distribution of Physiologic Indices.**



Distributions of (A) fractional flow reserve, (B) coronary flow reserve, (C) index of microcirculatory resistance, (D) resistive reserve ratio are shown.


**Figure S2. Correlation of Resistive Reserve Ratio with Diameter Stenosis or Other Physiologic Indices.**



Correlation of RRR with **(A)** percent diameter stenosis, **(B)** FFR, **(C)** CFR, and **(D)** IMR.


CFR, coronary flow reserve; FFR, fractional flow reserve; IMR, index of microcirculatory resistance; %DS, percent diameter stenosis; RRR, resistive reserve ratio.

**Figure S3. Distribution of CFR, FFR and Resistive Reserve Ratio.**



Distribution of vessels according to CFR and FFR values. Red dots represented vessels with depressed resistive reserve ratio (RRR)  $< 3.5$  and blue dots represent those with preserved RRR  $\geq 3.5$ . Although RRR showed high correlation with CFR ( $R=0.948$ ,  $p<0.001$ ), the classification agreement between CFR and RRR was only modest (Kappa value 0.605,  $p<0.001$ ) and 28.6% of patients with CFR  $> 2.0$  showed low RRR ( $< 3.5$ ). Numbers represent the number of vessels in each quadrant.

**Figure S4. Comparison of Patient-Oriented Composite Outcome According to Resistive Reserve Ratio.**



Cumulative incidence of patient-oriented composite outcomes (POCO) is shown according to cut-off value of resistive reserve ratio (RRR). Unadjusted hazard ratio and 95% confidence intervals are presented. CI, confidence intervals; HR, hazard ratios; RRR, resistive reserve ratio.